



Park Tree Inventory Summary Report

City of Washington, Missouri

April 2018

Prepared for:
City of Washington
Parks Department
1220 S Lakeshore Drive
Washington, Missouri 63090

Prepared by:
Davey Resource Group, Inc.
1500 North Mantua Street
Kent, Ohio 44240
800-828-8312



Acknowledgments

The City of Washington's vision to promote and preserve the urban forest and improve the management of public trees was a fundamental inspiration for this project. This vision will ensure canopy continuity, which will reduce stormwater runoff and improve air quality, public health, and aesthetic values.

The City of Washington is thankful for the grant funding it received from Missouri Department of Conservation (MDC) in cooperation with the Missouri Community Forestry Council and the U.S. Forest Service through its Tree Resource Improvement and Maintenance (TRIM) cost-share program. The TRIM grant program is designed to encourage communities to create and support long-term and sustained urban and community forestry programs throughout Missouri.



Notice of Disclaimer: Inventory data provided by Davey Resource Group, Inc. "DRG" are based on visual recording at the time of inspection. Visual records do not include individual testing or analysis, nor do they include aerial or subterranean inspection. DRG is not responsible for the discovery or identification of hidden or otherwise non-observable hazards. Records may not remain accurate after inspection due to the variability of inventoried material. DRG provides no warranty with respect to the fitness of the urban forest for any use or purpose whatsoever. Clients may choose to accept or disregard DRG's recommendations, or to seek additional advice. Important: know and understand that visual inspection is confined to the designated subject tree(s) and that the inspections for this project are performed in the interest of facts of the tree(s) without prejudice to or for any other service or any interested party.

Executive Summary

The City of Washington Parks Department commissioned an inventory and assessment of trees located within the parks and public facilities. Understanding an urban forest's structure, function, and value can promote management decisions that will improve public health and environmental quality. DRG collected and analyzed the inventory data to understand species composition and tree condition and to generate maintenance recommendations. Tree values and benefits have been quantified using the i-Tree Streets benefits model (developed by the United States Department of Agriculture Forest Service in partnership with The Davey Tree Expert Company). This report will discuss the health and benefits of the inventoried street tree population throughout all the parks of the City of Washington.

Key Findings

- The overall condition of the tree population is Good.
- The most common species are: *Quercus palustris* (pin oak), 7%; *Acer rubrum* (red maple), 5%; *Cercis canadensis* (eastern redbud), 5%; *Platanus occidentalis* (American sycamore), 5%; and *Q. bicolor* (swamp white oak), 5%.
- The vast majority (61%) of the urban forest is in the young, 0–8 inches DBH class.
- Approximately 49% of the population is recommended for a Young Tree Training Prune, 48% is recommended for a Tree Clean, and 3% is recommended for Removal.
- The park's tree population provides approximately \$138,184.05 in the following annual benefits:
 - *Aesthetic and Other Tangible Benefits*: valued at \$89,067.03 per year.
 - *Air Quality*: valued at \$3,556.52 per year.
 - *Net Total Carbon Sequestered and Avoided*: 579,559 pounds valued at \$1,655.75 per year.
 - *Energy*: 121.1 megawatt-hours (MWh) and 3,242 therms valued at \$11,390.50 per year.
 - *Stormwater*: 5,244,097 gallons valued at \$32,513.4 per year.

Table of Contents

Acknowledgments	i
Executive Summary	ii
Section 1: Tree Inventory Assessment	1
Section 2: i-Tree Streets Benefits	6
Conclusion and Recommendations.....	8
Glossary	9
References.....	13

Tables

1. Annual Benefits Provided by Inventoried Tree Population.....	6
2. Annual Energy Savings Provided by Inventoried Tree Population.....	6
3. Annual Air Quality Benefits Provided by Inventoried Tree Population	7
4. Annual Carbon Dioxide Benefits Provided by Inventoried Tree Population	7
5. Annual Stormwater Benefits Provided by Inventoried Tree Population	7
6. Annual Aesthetic and Other Tangible Benefits Provided by Inventoried Tree Population	7

Figures

1. Tree species composition in all Washington, Missouri parks	1
2. Top five genera across all Washington, Missouri parks in relation to the 20% Rule.	2
3. Age class distribution compared to Richards' (1983) ideal.	3
4. Overall condition of the population.....	3
5. Tree condition by age class.	4
6. Maintenance needs by risk rating.	5

Appendices

- A. Site Location Methods
- B. Tree Inventory Analysis Reports

Section 1: Tree Inventory Assessment

Project Area

In February 2018, DRG arborists assessed and inventoried trees in the parks and public facilities of the City of Washington. See Appendix A for an overview of the site location methodology used in the inventory and assessment.

Species Diversity

Throughout Washington's parks, 2,707 trees were inventoried. Figure 1 shows the composition of the most populous species compared to all inventoried species. The composition of a tree population should follow the 10-20-30 Rule for species diversity: a single species should represent no more than 10% of the urban forest, a single genus no more than 20%, and a single family no more than 30%.

Of the species found within the parks of Washington, no single species exceeds the 10% threshold. *Quercus palustris* (pin oak) is the closest to the threshold, comprising 7% of the total inventoried population.




Figure 1. Tree species composition in all Washington, Missouri parks.

Figure 2 compares the percentages of the most common genera identified during the inventory to the 20% Rule. *Quercus* (oak) exceeds the recommended 20% threshold for a single genus in a population. Oak comprises 24% of the park tree population.




Figure 2. Top five genera across all Washington, Missouri parks in relation to the 20% Rule.

Diameter Size Class Distribution

Analyzing the diameter size class distribution (measured as diameter at breast height [DBH]) provides an estimate of the relative age of a tree population and lends insight into maintenance practices and needs.

The inventoried trees were categorized into the following diameter size classes: young trees (0–8 inches DBH); established trees (9–17 inches DBH); maturing trees (18–24 inches DBH); and mature trees (greater than 24 inches DBH). These categories were chosen so that the population could be analyzed following Richards' ideal distribution (1983). Richards proposed an ideal diameter size class distribution for park trees based on observations of well-adapted trees in Syracuse, New York. Richards' ideal distribution suggests that the largest fraction of trees (approximately 40% of the population) should be young (less than 8 inches DBH), while a smaller fraction (approximately 10%) should fall in the large-diameter size class (greater than 24 inches DBH). A tree population with an ideal distribution would have an abundance of newly planted and young trees, and lower numbers of established, maturing, and mature trees.

Figure 3 compares Washington's inventoried park tree diameter size class distribution to the ideal proposed by Richards (1983). Washington Park's distribution trends toward the ideal; however, established and maturing trees fall short of the ideal by nearly 14% and 9%, respectively. As the urban forest in Washington ages, this ideal will begin to balance out. Continued tree planting, care, and maintenance of the young tree population will help achieve a more sustainable size distribution of park trees in Washington.




Figure 3. Age class distribution compared to Richards' (1983) ideal.

Condition

Several factors were considered for the condition of each tree, including: root characteristics, branch structure, trunk, canopy, foliage condition, and the presence of pests. The condition of each inventoried tree was rated Excellent, Good, Fair, Poor, Critical, or Dead.

Most of the inventoried Park trees were recorded to be in Good or Fair condition, 71% and 22%, respectively (Figure 4). Based on these data, the general health of the overall inventoried tree population is rated Good.




Figure 4. Overall condition of the population.

Figure 5 illustrates the general condition of the urban forest in relation to the relative age classes. The majority of young, established, maturing, and mature trees were rated to be in Good condition. With proactive care and an established maintenance schedule, the parks can improve the long-term health of its urban forest.




Figure 5. Tree condition by age class.

Primary Maintenance and Risk

Primary maintenance refers to the task identified for a tree or site: Removal, Tree Clean, or Young Tree Train. Risk is a graduated scale that measures potential tree-related hazardous conditions. A tree is considered hazardous when its potential risks exceed an acceptable level.

DRG based the maintenance recommendations and risk values (Figure 6) on the evaluation of species, diameter class, condition, impact of hazard, and defects found in each individual tree. Identifying and ranking the maintenance needs of a tree population enable tree work to be assigned priority based on observed defects. Once prioritized, tree work can be systematically addressed to eliminate the greatest risk and liability first (Stamen 2011).

Based on the inventoried population across all parks, the following maintenance recommendations should be implemented: 84 removals, 214 Tree Prunes, and 1,322 Young Tree Trains. There were 1,076 trees that were listed as Discretionary Prune. These trees should be taken care of after the Removals and Prunes. Figure 6 illustrates the risk values associated with each maintenance need.




Figure 6. Maintenance needs by risk rating.

Section 2: i-Tree Streets Benefits

The i-Tree Streets application was used to assess the inventoried trees. This management and analysis tool uses tree inventory data to quantify the dollar value of annual environmental and aesthetic benefits provided by trees, including energy conservation, air quality improvement, carbon dioxide (CO₂) reduction, stormwater control, and increases in property value. The tool estimates the costs and benefits of a park tree population and creates annual benefit reports that reflect the value park trees provide to a community.

The inventoried urban forest of Washington provides an annual benefit of approximately \$138,184 in energy savings, overall air quality improvements, carbon dioxide removal, stormwater reduction, and aesthetic and other tangible benefits. More detailed information about annual benefit analysis reports and maintenance reports can be found in Appendix B.

Figure 7 provides a breakdown of the annual benefits provided to the Parks and the City of Washington.

Table 1. Annual Benefits Provided by Inventoried Tree Population

Benefits	Total (\$)	Percent of Total
Aesthetics/Other	\$89,067.88	64%
Air Quality	\$3,556.52	3%
Carbon Dioxide	\$1,655.75	1%
Energy	\$11,390.50	8%
Stormwater	\$32,513.40	24%
All Park Trees Total	\$138,184.05	100%

Trees and Energy Use

Public trees conserve energy by shading structures and surfaces, reducing electricity use for air conditioning in the summer, and diverting wind in the winter to reduce natural gas use. Based on the inventoried trees, the annual electric and natural gas savings are equivalent to 121.1 MWh of electricity and 3,241.8 therms of natural gas. When converted into monetary values using default economic data, this accounts for an annual savings of \$11,390.50 in energy consumption. Large leafy canopies provide greater reductions in energy use by providing shade and natural wind barriers. In contrast, smaller trees tend to result in smaller reductions in energy usage.

Table 2. Annual Energy Savings Provided by Inventoried Tree Population

Total Electricity (MWh)	Total Natural Gas (Therms)	Total (\$)	Avg. \$/Tree
121.1	3,241.8	\$11,390.50	\$4.21

Air Quality Improvements

The inventoried tree population annually removes 2,406 pounds of air pollutants (including ozone, nitrogen dioxide, sulfur dioxide, and particulate matter) through deposition. The specified park tree population provides a total annual benefit of \$3,556.52 in overall air quality improvements. Each tree provides an average of \$1.31 in annual benefits.

Table 3. Annual Air Quality Benefits Provided by Inventoried Tree Population

Net Total Avoided/Deposition (lbs)	Total (\$)	Avg. \$/tree
2,406	\$3,556.52	\$1.31

Carbon Dioxide Removal

Trees store some of the CO₂ they absorb, thereby preventing it from reaching the upper atmosphere where it can react with other compounds and form harmful gases like ozone, which adversely affect air quality.

Through sequestration and avoidance, 579,559 pounds of CO₂ are removed on an annual basis. The inventoried trees provide an annual carbon benefit of \$1,655.75, with a per-tree average of \$0.61.

Table 4. Annual Carbon Dioxide Benefits Provided by Inventoried Tree Population

Net Total CO ₂ Sequestered/Avoided - Released (lbs.)	Total (\$)	Avg. \$/tree
579,559	\$1,655.75	\$0.61

Stormwater Interception and Mitigation

Trees intercept rainfall, which reduces costs to manage stormwater runoff. Washington's inventoried park trees intercept 5,244,097 gallons of rainfall annually. The estimated average savings for the parks and the City of Washington in stormwater management is approximately \$32,513.40 per year. On average, each tree provides an annual stormwater benefit of \$12.01.

Table 5. Annual Stormwater Benefits Provided by Inventoried Tree Population

Total Rainfall Interception (gal.)	Total (\$)	Avg. \$/tree
5,244,097	\$32,513.40	\$12.01

Aesthetic/Other Benefits

The total annual benefit associated with property value increases and other tangible and intangible services trees provide is \$89,067.99. Each tree provided an average benefit of \$32.90.

Table 6. Annual Aesthetic and Other Tangible Benefits Provided by Inventoried Tree Population

Total (\$)	Avg. \$/Tree
\$89,067.99	\$32.90

Conclusion and Recommendations

Managing trees in urban areas is often complicated. Navigating the recommendations of experts, the needs of residents, the pressures of local economics and politics, concerns for public safety and liability, physical components of trees, forces of nature and severe weather events, and the expectation that these issues are resolved all at once is a considerable challenge.

The parks and the City of Washington must carefully consider these challenges to fully understand the needs of maintaining an urban forest. By completing a tree inventory, the city has shown interest in preserving the urban forest, but also maintaining it for future generations. If the city successfully implements established planting and maintenance programs that include Young Tree Training, Routine Pruning, and public outreach, the health and safety of Washington's trees and residents will be maintained for years to come.

Washington's urban forest is in Good condition and provides \$138,184 in annual benefits. With continued dedication to its park tree resource, the city can improve the condition and diversity of its trees and increase the annual benefits they provide.

Glossary

Aesthetic/Other Report: The i-Tree Streets Aesthetic/Other Report presents the tangible and intangible benefits of trees reflected in increases in property values in dollars (\$).

Air Quality Report: The i-Tree Streets Air Quality Report quantifies the air pollutants (ozone [O₃], nitrogen dioxide [NO₂], sulfur dioxide [SO₂], coarse particulate matter less than 10 micrometers in diameter [PM₁₀]) deposited on tree surfaces, and reduced emissions from power plants (NO₂, PM₁₀, Volatile Oxygen Compounds [VOCs], SO₂) due to reduced electricity use measured in pounds (lbs.). Also reported are the potential negative effects of trees on air quality due to Biogenic Volatile Organic Compounds (BVOC) emissions.

arboriculture: The art, science, technology, and business of commercial, public, and utility tree care.

canopy: Branches and foliage that make up a tree's crown.

Carbon Dioxide Report: The i-Tree Streets Carbon Dioxide Report presents annual reductions in atmospheric CO₂ due to sequestration by trees and reduced emissions from power plants due to reduced energy use in pounds. The model accounts for CO₂ released as trees die and decompose and CO₂ released during the care and maintenance of trees.

clean (primary maintenance need): Based on *ANSI A300 (Part 1)* standards, selective removal of dead, dying, broken, and/or diseased wood to minimize potential risk.

community forest: see **urban forest**.

condition (data field): The general condition of each tree rated during the inventory according to the following categories adapted from the International Society of Arboriculture's rating system: Excellent (100%), Very Good (90%), Good (80%), Fair (60%), Poor, (40%), Critical (20%), Dead (0%).

diameter at breast height (DBH): See **tree size**.

diameter: See **tree size**.

Energy Report: The i-Tree Streets Energy Report presents the contribution of the urban forest toward conserving energy in terms of reduced natural gas use in winter measured in therms [th] and reduced electricity use for air conditioning in summer measured in megawatt-hours (MWh).

failure: In terms of tree management, failure is the breakage of stem or branches, or loss of mechanical support of the tree's root system.

genus: A taxonomic category ranking below a family and above a species and generally consisting of a group of species exhibiting similar characteristics. In taxonomic nomenclature, the genus name is used, either alone or followed by a Latin adjective or epithet, to form the name of a species.

geographic information system (GIS): A technology that is used to view and analyze data from a geographic perspective. The technology is a piece of an organization's overall information system framework. GIS links location to information (such as people to addresses, buildings to parcels, or streets within a network) and layers that information to give you a better understanding of how it all interrelates.

global positioning system (GPS): GPS is a system of earth-orbiting satellites that makes it possible for people with ground receivers to pinpoint their geographic location.

High Risk tree: Tree that cannot be cost-effectively or practically treated. Most High Risk trees have multiple or significant defects affecting less than 40% of the trunk, crown, or critical root zone. Defective trees and/or tree parts are most likely between 4–20 inches in diameter and can be found in areas of frequent occupation, such as a main thoroughfare, congested streets, and/or near schools.

Importance Values: A calculation in i-Tree Streets. Importance Values (IV) are displayed in table form for all species that make up more than 1% of the population. The Streets IV is the mean of three relative values (percentage of total trees, percentage of total leaf area, and percentage of canopy cover) and can range from 0 to 100 with an IV of 100 suggesting total reliance on one species. IVs offer valuable information about a community's reliance on certain species to provide functional benefits. For example, a species might represent 10% of a population, but have an IV of 25% because of its great size, indicating that the loss of those trees due to pests or disease would be more significant than their numbers suggest.

inventory: See **tree inventory**.

i-Tree Streets: i-Tree Streets is a street tree management and analysis tool that uses tree inventory data to quantify the dollar value of annual environmental and aesthetic benefits: energy conservation, air quality improvement, CO₂ reduction, stormwater control, and property value increase.

i-Tree Tools: State-of-the-art, peer-reviewed software suite from the USDA Forest Service that provides urban forestry analysis and benefits assessment tools. The i-Tree Tools help communities of all sizes to strengthen their urban forest management and advocacy efforts by quantifying the structure of community trees and the environmental services that trees provide.

Low Risk tree: Tree with minor visible structural defects or wounds in areas with moderate to low public access.

mapping coordinate (data field): Helps to locate a tree; X and Y coordinates were generated for each tree using GPS.

Moderate Risk tree: Tree with defects that may be cost-effectively or practically treated. Most of the trees in this category exhibit several moderate defects affecting more than 40% of a tree's trunk, crown, or critical root zone.

monoculture: A population dominated by one single species or very few species.

Net Annual Benefits: Specific data field for i-Tree Streets. Citywide benefits and costs are calculated according to category and summed. Net benefits are calculated as benefits minus costs.

Nitrogen Dioxide (NO₂): Nitrogen dioxide is a compound typically created during the combustion processes and is a major contributor to smog formation and acid deposition.

Ozone (O₃): A strong-smelling, pale blue, reactive toxic chemical gas with molecules of three oxygen atoms. It is a product of the photochemical process involving the Sun's energy. Ozone exists in the upper layer of the atmosphere as well as at the Earth's surface. Ozone at the Earth's surface can cause numerous adverse human health effects. It is a major component of smog.

Particulate Matter (PM₁₀): A major class of air pollutants consisting of tiny solid or liquid particles of soot, dust, smoke, fumes, and mists.

primary maintenance need (data field): The type of tree work needed to reduce immediate risk.

pruning: The selective removal of plant parts to meet specific goals and objectives.

removal (primary maintenance need): Data field collected during the inventory identifying the need to remove a tree. Trees designated for removal have defects that cannot be cost-effectively or practically treated. Most of the trees in this category have a large percentage of dead crown.

right-of-way (ROW): See **street right-of-way**.

risk: Combination of the probability of an event occurring and its consequence.

risk assessment (data fields): The risk assessment is a point-based assessment of each tree by an arborist using a protocol based on the U.S. Forest Service Community Tree Risk Rating System. In the field, the probability of tree or tree part failure is assigned 1–4 points (identifies the most likely failure and rates the likelihood that the structural defect(s) will result in failure based on observed, current conditions); the size of defective tree part is assigned 1–3 points (rates the size of the part most likely to fail); the probability of target impact by the tree or tree part is assigned 1–3 points (rates the use and occupancy of the area that would be struck by the defective part); and other risk factors are assigned 0–2 points (used if professional judgment suggests the need to increase the risk rating). The data from the risk assessment is used to calculate the risk rating that is ultimately assigned to the tree.

risk rating: Calculated from the field risk assessment data (see **risk assessment**), this is the sum of total risk assessment values. Risk ratings range from 3–10, with 3 being the lowest risk and 10 being the highest risk. In this Plan, the risk rating was used to identify the severity of risk assigned to a tree and to prioritize tree maintenance needs. The following categories were used:

- risk rating of 9 or 10 = Severe Risk tree
- risk rating of 7 or 8 = High Risk tree
- risk rating of 5 or 6 = Moderate Risk tree
- risk rating of 3 or 4 = Low Risk tree
- risk rating of 0 = no risk (used only for planting sites and stumps)

secondary maintenance need (data field): Recommended maintenance for a tree, which may be risk oriented, such as raising the crown for clearance, but generally was geared toward improving the structure of the tree and enhancing aesthetics.

Severe Risk tree: Tree rated to be Severe Risk cannot be cost-effectively or practically treated. Most Severe Risk trees have multiple and significant defects present in the trunk, crown, or critical root zone. Defective trees and/or tree parts are most likely larger than 20 inches in diameter and can be found in areas of frequent occupation, such as a main thoroughfare, congested streets, and/or near schools.

species: Fundamental category of taxonomic classification, ranking below a genus or subgenus, and consisting of related organisms capable of interbreeding.

stem: A woody structure bearing buds and foliage, and giving rise to other stems.

stems (data field): Identifies the number of stems or trunks splitting less than 1 foot above ground level.

Stored Carbon Report: Whereas, the i-Tree Streets Carbon Dioxide Report quantifies annual CO₂ reductions, and the i-Tree Streets Stored Carbon Report tallies all of the Carbon (C) stored in the urban forest over the life of the trees as a result of sequestration measured in pounds as the CO₂ equivalent.

Stormwater Report: A report generated by i-Tree Streets that presents the reductions in annual stormwater runoff due to rainfall interception by trees measured in gallons (gals.).

street name (data field): The name of a street right-of-way or road identified using posted signage or parcel information.

street right-of-way (ROW): A strip of land generally owned by a public entity over which facilities, such as highways, railroads, or power lines, are built.

street tree: A street tree is defined as a tree within the right-of-way.

structural defect: A feature, condition, or deformity of a tree or tree part that indicates weak structure and contributes to the likelihood of failure.

stump removal (primary maintenance need): Indicates a stump that should be removed.

Sulfur Dioxide (SO₂): A strong-smelling, colorless gas that is formed by the combustion of fossil fuels. Sulfur oxides contribute to the problem of acid rain.

Summary Report: The i-Tree Streets Summary report presents the annual total of energy, stormwater, air quality, carbon dioxide, and aesthetic/other benefits. Values are dollars per tree or total dollars.

tree benefit: An economic, environmental, or social improvement that benefits the community and results mainly from the presence of a tree. The benefit received has real or intrinsic value associated with it.

tree inventory: Comprehensive database containing information or records about individual trees typically collected by an arborist.

tree size (data field): A tree's diameter measured to the nearest inch in 1-inch size classes at 4.5 feet above ground, also known as diameter at breast height (DBH) or diameter.

tree: A tree is defined as a perennial woody plant that may grow more than 20 feet tall. Characteristically, it has one main stem, although many species may grow as multi-stemmed forms.

urban forest: All of the trees within a municipality or a community. This can include the trees along streets or rights-of-way, in parks and greenspaces, in forests, and on private property.

Volatile Organic Compounds (VOCs): Hydrocarbon compounds that exist in the ambient air and are by-products of energy used to heat and cool buildings. Volatile organic compounds contribute to the formation of smog and/or are toxic. Examples of VOCs are gasoline, alcohol, and solvents used in paints.

Young Tree Train (primary maintenance need): Data field based on *ANSI A300 (Part 1)* standards, pruning of young trees to correct or eliminate weak, interfering, or objectionable branches to improve structure. These trees, up to 20 feet in height, can be worked with a pole pruner by a person standing on the ground.

References

- Richards, N.A. 1983. "Diversity and Stability in a Street Tree Population." *Urban Ecology* 7(2):159–171.
- Stamen, R.S. "Understanding and Preventing Arboriculture Lawsuits." Presented at the Georgia Urban Forest Council Annual Meeting, Madison, Georgia, November 2–3, 2011.

Appendix A

Data Collection and Site Location Methods

Data Collection Methods

DRG collected tree inventory data using a system that utilizes a customized ArcPad program loaded onto pen-based field computers equipped with geographic information system (GIS) and global positioning system (GPS) receivers. The knowledge and professional judgment of Davey Resource Group's arborists ensure the high quality of inventory data.

Data fields are defined in the glossary of the Summary Report. At each site, the following data fields were collected:

- defects
- condition
- further inspection
- primary maintenance needs
- mapping coordinates
- multi-stem
- park or property name
- observations
- overhead utilities
- notes
- residual risk
- risk rating
- species
- tree size*

* measured in inches in diameter at 4.5 feet above ground (or diameter at breast height [DBH])

Maintenance needs are based on *ANSI A300 (Part 1)* (ANSI 2008). Risk assessment and risk rating are based on *Best Management Practices: Tree Risk Assessment* (International Society of Arboriculture [ISA] 2011).

Site Location Methods

Equipment and Base Maps

Inventory arborists use FZ-G1 Panasonic Toughpad® unit(s).

Base map layers were loaded onto these unit(s) to help locate sites during the inventory. Table 1 lists the base map layers, utilized along with source and format information for each layer.

Table 1. Base Map Layers Utilized for Inventory

Imagery/Data Source	Date	Projection
Missouri Spatial Data Clearinghouse http://www.msdis.missouri.edu/data/themelist.html Aerial Imagery is from 2012	2012 & 2016	NAD 1983 UTM Zone 15N, Meters

Park and/or Public Space Site Location

Park and/or public space site locations were collected using the park and/or public space's name.

Appendix B
Tree Inventory Analysis Reports

Matrix Report

Park Name by Primary Maintenance Need

	Removal	Discretionary	Prune	Train	Total
Bernie E. Hillermann Park	15	414	66	373	868
Burger Park	3	37	14	142	196
James W. Rennick Riverfront Park	10	43	26	44	123
Krog Memorial Park	2	48	10	21	81
Lafayette Plaza	0	2	0	0	2
Jerry J. Jasper Lakeview Park	17	181	19	112	329
McLaughlin Field	0	2	1	22	25
Optimist Park	3	44	12	19	78
Phoenix Park	0	65	0	210	275
Rotary Riverfront Trail	19	45	38	132	234
Washington City Park	5	60	19	21	105
Crestview Water Tower	2	20	0	51	73
Eckelcamp Park	0	14	1	9	24
City Hall	0	2	1	78	81
Farmers Market	0	0	0	9	9
Train Station	0	0	0	6	6
Third Parkway	1	14	0	4	19
First Parkway	1	11	6	1	19
9th Street Commons	2	22	3	4	31
South Point Bark Park	1	18	6	14	39
Jefferson Street	0	0	0	16	16
None	0	3	0	0	3
Fire Stations	2	17	0	17	36
Water Treatment Facility	1	6	2	1	10
Well House	0	2	1	5	8
Parking Lot	0	6	0	11	17

Total	84	1076	225	1322	2707
--------------	-----------	-------------	------------	-------------	-------------

**The headings and numbers act as links to search the sites from the results of the matrix.*

Washington, MO

TreeKeeper 7

Matrix Report

Park Name by Risk Rating

	High	Low	Moderate	Total
Bernie E. Hillermann Park	0	860	8	868
Burger Park	0	196	0	196
James W. Rennick Riverfront Park	1	121	1	123
Krog Memorial Park	0	76	5	81
Lafayette Plaza	0	1	1	2
Jerry J. Jasper Lakeview Park	0	329	0	329
McLaughlin Field	0	25	0	25
Optimist Park	0	78	0	78
Phoenix Park	0	275	0	275
Rotary Riverfront Trail	0	231	3	234
Washington City Park	1	101	3	105
Crestview Water Tower	0	73	0	73
Eckelcamp Park	0	24	0	24
City Hall	0	81	0	81
Farmers Market	0	9	0	9
Train Station	0	6	0	6
Third Parkway	0	19	0	19
First Parkway	0	18	1	19
9th Street Commons	0	30	1	31
South Point Bark Park	0	38	1	39
Jefferson Street	0	16	0	16
None	0	3	0	3
Fire Stations	0	36	0	36
Water Treatment Facility	0	10	0	10
Well House	0	8	0	8
Parking Lot	0	17	0	17

Total

2 2681 24 2707

**The headings and numbers act as links to search the sites from the results of the matrix.*

Washington, MO

TreeKeeper 7

Tree Frequency Report

<i>Click for locations</i>	<i>Count</i>	<i>Percentage(%)</i>
<u>Quercus palustris (oak, pin)</u>	177	6.54
<u>Platanus occidentalis (sycamore, American)</u>	148	5.47
<u>Acer rubrum (maple, red)</u>	144	5.32
<u>Cercis canadensis (redbud, eastern)</u>	132	4.88
<u>Quercus bicolor (oak, swamp white)</u>	129	4.77
<u>Taxodium distichum (baldcypress, common)</u>	124	4.58
<u>Quercus macrocarpa (oak, bur)</u>	109	4.03
<u>Populus deltoides (cottonwood, eastern)</u>	86	3.18
<u>Betula nigra (birch, river)</u>	82	3.03
<u>Quercus muehlenbergii (oak, chinkapin)</u>	74	2.73
<u>Acer negundo (boxelder)</u>	70	2.59
<u>Ulmus pumila (elm, Siberian)</u>	66	2.44
<u>Acer saccharum (maple, sugar)</u>	60	2.22
<u>Malus spp. (crabapple, flowering)</u>	59	2.18
<u>Juglans nigra (walnut, black)</u>	59	2.18
<u>Quercus rubra (oak, northern red)</u>	56	2.07
<u>Acer saccharinum (maple, silver)</u>	55	2.03
<u>Ulmus americana (elm, American)</u>	54	1.99
<u>Fraxinus pennsylvanica (ash, green)</u>	53	1.96
<u>Pinus taeda (pine, loblolly)</u>	47	1.74

<u>Celtis occidentalis (hackberry, common)</u>	47	1.74
<u>Thuja occidentalis (arborvitae, eastern)</u>	44	1.63
<u>Pinus strobus (pine, eastern white)</u>	44	1.63
<u>Pyrus calleryana (pear, callery)</u>	44	1.63
<u>Quercus alba (oak, white)</u>	42	1.55
<u>Fraxinus americana (ash, white)</u>	41	1.51
<u>Quercus imbricaria (oak, shingle)</u>	34	1.26
<u>Cornus racemosa (dogwood, gray)</u>	33	1.22
<u>Nyssa sylvatica (blackgum)</u>	31	1.15
<u>Lagerstroemia indica (crapemyrtle, common)</u>	30	1.11
<u>Liriodendron tulipifera (tuliptree)</u>	25	0.92
<u>Gymnocladus dioica (Kentucky coffeetree)</u>	24	0.89
<u>Gleditsia triacanthos inermis (honeylocust, thornless)</u>	24	0.89
<u>Cornus florida (dogwood, flowering)</u>	23	0.85
<u>Ginkgo biloba (ginkgo)</u>	23	0.85
<u>Amelanchier spp. (serviceberry, spp.)</u>	20	0.74
<u>Morus alba (mulberry, white)</u>	18	0.66
<u>Cornus kousa (dogwood, Kousa)</u>	17	0.63
<u>Prunus serotina (cherry, black)</u>	16	0.59
<u>Prunus spp. (cherry/plum, spp.)</u>	16	0.59
<u>Salix nigra (willow, black)</u>	15	0.55
<u>Quercus phellos (oak, willow)</u>	11	0.41

<u>Juniperus virginiana</u> (redcedar, eastern)	11	0.41
<u>Catalpa speciosa</u> (catalpa, northern)	10	0.37
<u>Viburnum rufidulum</u> (viburnum, rusty blackhaw)	10	0.37
<u>Sassafras albidum</u> (sassafras)	10	0.37
<u>Acer platanoides</u> (maple, Norway)	10	0.37
<u>Viburnum lentago</u> (nannyberry)	10	0.37
<u>Zelkova serrata</u> (zelkova, Japanese)	10	0.37
<u>Ulmus parvifolia</u> (elm, Chinese)	10	0.37
<u>Quercus robur</u> (oak, English)	9	0.33
<u>Acer x freemanii</u> (maple, Freeman)	9	0.33
<u>Liquidambar styraciflua</u> (sweetgum, American)	9	0.33
<u>Diospyros virginiana</u> (persimmon, common)	9	0.33
<u>Aesculus glabra</u> (buckeye, Ohio)	8	0.30
<u>Acer palmatum</u> (maple, Japanese)	8	0.30
<u>Pinus nigra</u> (pine, Austrian)	7	0.26
<u>Carya illinoensis</u> (pecan)	7	0.26
<u>Robinia pseudoacacia</u> (locust, black)	7	0.26
<u>Tsuga canadensis</u> (hemlock, eastern)	7	0.26
<u>Ilex opaca</u> (holly, American)	7	0.26
<u>Tilia cordata</u> (linden, littleleaf)	7	0.26
<u>Ilex decidua</u> (possumhaw)	5	0.18
<u>Pinus echinata</u> (pine, shortleaf)	5	0.18

<u>Acer campestre (maple, hedge)</u>	5	0.18
<u>Picea pungens (spruce, Colorado)</u>	5	0.18
<u>Chionanthus virginicus (fringetree, white)</u>	5	0.18
<u>Acer tataricum ginnala (maple, Amur)</u>	5	0.18
<u>Carya cordiformis (hickory, bitternut)</u>	5	0.18
<u>Rhus typhina (sumac, staghorn)</u>	5	0.18
<u>Carya glabra (hickory, pignut)</u>	5	0.18
<u>Ulmus x (elm, hybrid)</u>	4	0.15
<u>Hamamelis virginiana (witchhazel, common)</u>	4	0.15
<u>Carya tomentosa (hickory, mockernut)</u>	4	0.15
<u>Taxodium ascendens (pondcypress)</u>	4	0.15
<u>Asimina triloba (pawpaw, common)</u>	3	0.11
<u>Gleditsia triacanthos (honeylocust)</u>	3	0.11
<u>Magnolia x soulangiana (magnolia, saucer)</u>	3	0.11
<u>Koelreuteria paniculata (goldenraintree)</u>	3	0.11
<u>Castanea mollissima (chestnut, Chinese)</u>	2	0.07
<u>Fagus grandifolia (beech, American)</u>	2	0.07
<u>Crataegus spp. (hawthorn, spp.)</u>	2	0.07
<u>Picea abies (spruce, Norway)</u>	2	0.07
<u>Carpinus caroliniana (hornbeam, American)</u>	2	0.07
<u>Quercus nigra (oak, water)</u>	2	0.07

<u>Halesia tetraptera (silverbell, Carolina)</u>	2	0.07
<u>Magnolia grandiflora (magnolia, southern)</u>	2	0.07
<u>Cladrastis kentukea (yellowwood)</u>	2	0.07
<u>Tilia americana (linden, American)</u>	2	0.07
<u>Ulmus rubra (elm, slippery)</u>	2	0.07
<u>Carya ovata (hickory, shagbark)</u>	2	0.07
<u>Eucommia ulmoides (hardy rubber tree)</u>	2	0.07
<u>Rhamnus caroliniana (buckthorn, Carolina)</u>	2	0.07
<u>Platanus x acerifolia (planetree, London)</u>	2	0.07
<u>Syringa reticulata (lilac, Japanese tree)</u>	2	0.07
<u>Aesculus pavia (buckeye, red)</u>	1	0.04
<u>Magnolia virginiana (magnolia, sweetbay)</u>	1	0.04
<u>Rhus copallinum (sumac, winged)</u>	1	0.04
<u>Cornus alternifolia (dogwood, pagoda)</u>	1	0.04
<u>Aesculus hippocastanum (horsechestnut)</u>	1	0.04
<u>Prunus americana (plum, American)</u>	1	0.04
<u>Magnolia acuminata (magnolia, cucumbertree)</u>	1	0.04
<u>Quercus acutissima (oak, sawtooth)</u>	1	0.04
<u>unknown tree (unknown tree)</u>	1	0.04
<u>Albizia julibrissin (mimosa)</u>	1	0.04

<u>Carya laciniosa (hickory, shellbark)</u>	1	0.04
<u>Picea rubens (spruce, red)</u>	1	0.04
<u>Euonymous atropurpurea (eastern wahoo)</u>	1	0.04
<u>Ostrya virginiana (hophornbeam, American)</u>	1	0.04
<u>Acer buergerianum (maple, trident)</u>	1	0.04
<u>Cornus mas (dogwood, corneliancherry)</u>	1	0.04
<u>Acer griseum (maple, paperbark)</u>	1	0.04
<u>Betula populifolia (birch, gray)</u>	1	0.04
<u>Pinus spp. (pine, spp.)</u>	1	0.04
<u>Rhus glabra (sumac, smooth)</u>	1	0.04
<u>Carpinus betulus (hornbeam, European)</u>	1	0.04

**2707 Species Found,
116 distinct species**

Washington, MO

TreeKeeper 7